



ELSEVIER

Are subtype differences important in HIV drug resistance?

RJ Lessells^{1,2}, DK Katzenstein³ and T de Oliveira^{1,4}

The diversity of human immunodeficiency virus type 1 (HIV-1) has given rise to multiple subtypes and recombinant strains. The majority of research into antiretroviral agents and drug resistance has been performed on subtype B viruses, yet non-subtype B strains are responsible for 90% of global infections. Although it seems that combination antiretroviral regimens are effective against all HIV-1 subtypes, there is emerging evidence of subtype differences in drug resistance, relevant to antiretroviral strategies in different parts of the world. For this purpose, extensive sampling of HIV genetic diversity, curation and analyses are required to inform antiretroviral strategies in different parts of the world.

Addresses

¹ Africa Centre for Health and Population Studies, University of KwaZulu-Natal, Somkhele, South Africa

² Department of Clinical Research, London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT, United Kingdom

³ Stanford University Medical Center, Department of Medicine, Stanford, CA 94305, USA

⁴ Research Department of Infection, University College of London, London, W1T 4JF, United Kingdom

Corresponding author: de Oliveira, T (tdeoliveira@africacentre.ac.za)

Current Opinion in Virology 2012, 2:636–643

This review comes from a themed issue on **Antivirals and resistance**

Edited by **Daniel Lamarre** and **Mark A Wainberg**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 21st September 2012

1879-6257/\$ – see front matter, © 2012 Elsevier B.V. All rights reserved.

<http://dx.doi.org/10.1016/j.coviro.2012.08.006>

Introduction

The past decade has seen substantial global scale-up of antiretroviral therapy (ART) for HIV infection and more than six million people are receiving ART in low-income and middle-income countries [1]. Antiretroviral drug resistance is one of the main threats to global control of HIV [2]. The majority of persons living with HIV infection are infected with non-subtype B variants of HIV type 1 (HIV-1) [3]. There is increasing evidence that polymorphisms that occur naturally in different HIV-1 subtypes impact on drug resistance and susceptibility to antiretroviral drugs.

Here, we outline the latest developments in subtyping tools, drug resistance databases and review recent evidence from *in vitro* and clinical studies regarding drug resistance among HIV-1 subtypes (Box 1).

HIV-1 origin, subtypes and recombinants

HIV-1 main group (group M) originated in West-Central Africa approximately 100 years ago [4,5^{••}]. It has since diversified into a large number of variants, including nine subtypes (A–D, F–H, J–K), six subsubtypes (A1–A4, F1–F2), multiple (>48) circulating recombinant forms (CRFs) and thousands of unique recombinant forms (URFs) (Los Alamos HIV Sequence Database; URL: <http://www.hiv.lanl.gov>) [5^{••},6]. The classification of recombinant viruses is based on complete genome analysis: CRFs are widespread, whereas URFs are restricted to a limited number of individuals [6]. The high number of existing HIV-1 variants is caused by both biological and epidemiological factors, which have been recently reviewed [4,5^{••},7].

HIV-1 variants are continually introduced into new populations by mobility and migration [3,5^{••},6,7]. As HIV-1 variants intermix in different part of the world, the likelihood of generating new recombinant viruses increases [6]. For example, a recent study in Quebec, Canada identified four subtypes, three CRFs and two new URFs. One of the new URFs is a recombinant of A/B (the RT/protease region was largely of subtype A, the integrase was subtype B), which is spreading and may be classified as a new CRF once complete genomes are sequenced [8]. Studies in London have detected all HIV-1 subtypes, the majority of CRFs and many previously undetected URFs [9,10]. Identification of individuals infected with different subtypes is increasing in metropolitan areas [8,11].

Subtyping tools and drug resistance databases

HIV-1 subtyping can be achieved by automated subtyping tools. At the time of this review, over 400 000 isolates have been subtyped using the Rega HIV-1 subtyping tool. This tool uses phylogenetic analysis to identify subtypes and CRFs. A recent upgrade has allowed the identification of many new CRFs and, for the first time, the classification of URFs [Rega HIV Subtyping Tool V3; URL: <http://www.bioafrica.net>]. Figure 1 shows a new feature of Rega Subtyping Tool V3, which is the phylogenetic identification of recombinant segments. A large comparison study of over 6000 sequences, carefully subtyped by phylogenetic methods, was conducted to evaluate the accuracy of REGAv3 and six other subtyping tools (ACP Pena *et al.* 17th International Bioinformatics Workshop on Virus Evolution and Molecular Epidemiology, Belgrade, Serbia, August 2012). The comparison tools included two new, sophisticated tools: SCUDEL [12] and COMET V2 (D Struck *et al.* 8th European HIV Drug Resistance Workshop, Sorrento, Italy, March 2010). The

Box 1 Summary of main concepts

- HIV-1 diversity has given rise to numerous subtypes and recombinant forms.
- New subtyping tools (e.g. Rega HIV-1 Subtyping Tool version 3, SCUEL and COMET) can accurately identify the most important HIV-1 variants.
- National and international public drug resistance databases are useful resources to trace the evolution of drug resistance in different subtypes.
- HIV-1 subtype genetic variation can influence the development of drug resistance and the susceptibility to certain antiretroviral drugs.
- K65R is an example of a clinically relevant mutation that emerges more frequently and more rapidly in subtype C viruses compared to subtype B; this has been shown to be related to the different template nucleotide sequence.
- Evidence from recent clinical trials and cohort studies suggests that response to combination antiretroviral regimens does not differ substantially by HIV-1 subtype.
- Appreciation of subtype differences is important in the development of new drugs and in the formulation of antiretroviral strategies.

three tools identified most of the pure subtypes in *pol* with high sensitivity and specificity (>95%). COMETv2 and REGAv3 identify the two most important CRFs (CRF01_AE and CRF02_AG) in more than 95%. Given that the great majority (>90%) of the infections in the world are owing to subtypes A, B C, CRF01_AE and CRF02_AG [3,5^{**},7], these recent subtyping tools can accurately identify most of the epidemiologically important HIV-1 variants and classify new recombinants.

International and country specific drug resistance databases are important repositories of HIV-1 genetic data [13]. The UK Drug Resistance Database contains over 10 000 non-B subtype isolates and proposals can be submitted for the use of data [UK HIV Drug Resistance Database; URL: <http://www.hivrd.org.uk>]. The Stanford HIV Drug Resistance Database (HIVDB) curates all published data and contains nearly 150 000 sequences. This is presented in many statistical and graphical formats [Stanford HIV Drug Resistance Database; URL: <http://hivdb.stanford.edu>]. HIVDB data and analyses have shown that, in spite of the large genetic variation found within subtypes, no major drug resistance mutation naturally occurs in naïve sequences [14,15]. However, a number of treatment-experienced mutation differences have been highlighted in the literature and are stored and curated in HIVDB. Table 1 summarises HIV drug resistance mutations associated with subtypes and CRFs in treatment experienced samples.

Nucleoside and nucleotide reverse transcriptase inhibitors


The lysine to arginine mutation at position 65 (K65R) is a major mutation that confers broad high-level resistance to most nucleoside/nucleotide reverse transcriptase inhibitors (NRTI/NtRTIs), except zidovudine. There is a substantial body of evidence that K65R emerges more frequently and more rapidly in subtype C viruses than in subtype B.

It has now been demonstrated that the difference in selection of K65R between subtypes B and C is related to the template nucleotide sequence and preferential pausing of reverse transcription at position 65. The nucleotide sequence at codons 64-65-66 differs between subtypes B and C and subtype C viruses contain a homopolymeric stretch of adenine bases. This leads to RT pausing during the synthesis of double-stranded DNA from the single-stranded DNA intermediate template, a process that is template-specific but independent of the RT enzyme [16]. Subsequent misalignment of the template and primer leads to the AAG to AGG change responsible for the K65R mutation [17^{*}].

Several recent studies have used ultra-deep pyrosequencing (UDPS) or allele-specific PCR (AS-PCR) to explore the frequency of low-level K65R mutants in both ART-naïve and ART-experienced individuals [18–22]. In one study using UDPS, the frequency of K65R at both >1% and >0.4% levels was higher in subtype C compared to subtype B and non-B/non-C subtypes [18]. In another study, the K65R mutation was detected at a higher frequency in ART-naïve subtype C individuals compared to those infected with subtype CRF01_AE (6% vs. 1%) and subtype B [19]. In ART-experienced patients with virological failure and without K65R on conventional Sanger sequencing, one study detected the presence of K65R by AS-PCR in 13% (4/30) of patients [20]; conversely other groups using UDPS detected no additional mutations in those without K65R on Sanger sequencing [21]. It is important to note the limitations of these highly sensitive sequencing techniques and spurious detection of the K65R mutation through PCR-induced mutation has been demonstrated [22].

There is recent clinical evidence demonstrating frequent and early emergence of K65R on tenofovir-based first-line ART regimens in South Africa [23^{*}]. Recent analysis of large scale implementation of TDF, 3TC and NVP indicates a higher rate of virological failure with this regimen [24^{*}]. In addition, a case report has documented the presence of low-level K65R mutants pre-treatment by clonal analysis, with enhancement of K65R variants within two months of treatment [25]. There is an urgent need for further research to determine the prevalence and impact of low-level K65R mutants, especially in settings where subtype C predominates and where tenofovir is now a component of first-line ART regimens.

Figure 1



Recombination profile and phylogenies of recombinant regions of a CRF03_AB isolate 03 AB,RU,97,KAL153 2 [Rega HIV Subtyping Tool V3; URL: <http://www.bioafrica.net>]. One of the new features of Rega Subtyping Tool version 3.0 is that it can perform detailed recombination analyses. The tool detects recombination, identifies the recombinant fragments and creates a phylogenetic tree fragments (Query sequence is at the top of the phylogenies). This figure shows a CRF recombinant A/B sequence (CRF03_AB, Genbank accession number AF193276). The subtype A region is from position 2252 to 2782 (Protease amino acid position 1–99 and RT 1–78) and subtype B from 2782 to 4822 (RT amino acid position 79–440 and Integrase amino acid position 1–198). Numbering is performed based on the complete genome reference sequence HXB2.

Non-nucleoside reverse transcriptase inhibitors

The emergence of non-nucleoside reverse transcriptase inhibitor (NNRTI) resistance mutation occurs after single dose nevirapine (sdNVP) for the prevention of mother-to-child HIV transmission. Previous work has suggested that this occurs more frequently with subtype C viruses [26,27]. One more recent study showed that

resistance mutations could be demonstrated using AS-PCR in 25% of patients more than 24 months after sdNVP exposure (C Yang *et al.* 17th Conference on Retroviruses and Opportunistic Infections, San Francisco, California, February 2010). However, the clinical significance of this is uncertain as the same study demonstrated no association between the presence of resistance mutations and virological failure on a subsequent NNRTI-based

Table 1

Recently described drug resistance mutations between HIV-1 subtypes and CRFs and impact on antiretroviral drug resistance and susceptibility

Position	Mutation	Comments	Reference
<i>Reverse transcriptase</i>			
RT65	K65R	Subtype C – AAG (K); subtype B – AAA (K): preferential pausing of reverse transcription, related to homopolymeric stretch of adenine bases	[16,17*]
RT138	E138K	E138K the first mutation to emerge in subtype C during etravirine therapy	[30]
RT181	Y181C	Preferential selection of Y181C for subtype A and B during etravirine therapy	[31]
RT348	N348I	Reduces susceptibility to etravirine in subtypes A, B and C. High prevalence in subtype C samples from patients failing first-generation NNRTIs	[32,33*]
<i>Protease</i>			
PR17	G17E	CRF2_AG hypersusceptibility to nelfinavir, atazanavir and indinavir	[38]
PR36	M36I	Subtype C – ATA (I); subtype B – ATG (M): affects susceptibility to protease inhibitors and viral replication capacity	[36]
PR64	I64M	CRF2_AG hypersusceptibility to nelfinavir, atazanavir and indinavir	[38]
PR89	M89T	Subtype C – ATG (M); subtype B – CTG (L): leads to preferential emergence of M89T in subtype C	[37]
<i>Integrase</i>			
IN92	E92Q	E92Q/N155H double mutant 10-fold more resistant to raltegravir and elvitegravir in subtype B versus subtype C	[47]
IN101	L101I	Present more frequently in non-B subtypes compared to subtype B (both INI-naïve and RAL-experienced)	[48]
IN118	G118R	Most common resistance pathway during dolutegravir therapy in subtype C	[50]
IN124	T124A	Present more frequently in INI-naïve non-B subtypes compared to subtype B	[48]
IN155	N155H	Subtype B with this mutation more resistant to raltegravir (and elvitegravir) than subtype C	[47]
IN263	R263K	Most common resistance pathway during dolutegravir therapy in subtype B	[49*]

regimen (PJ Weidle *et al.* 17th Conference on Retroviruses and Opportunistic Infections, San Francisco, California, February 2010).

Etravirine is a second-generation NNRTI which retains activity against strains with some resistance to nevirapine and efavirenz and which might therefore be an option as a component of a salvage regimen for antiretroviral-experienced patients. It has demonstrated good efficacy across subtypes [28]. There are a number of etravirine resistance-associated mutations (RAMs) which reduce the response to etravirine. These mutations are commonly present as polymorphisms in ART-naïve individuals infected with non-B subtypes, especially CRF02_AG [29]. There is conflicting evidence on the resistance pathways selected by etravirine therapy. One study found E138K the first mutation to emerge in subtypes B, C and CRF02_AG [30]. A separate study found the same for subtype C but demonstrated preferential selection of Y181C for subtype B virus [31].

A novel mutation in the C-terminal domain of RT (N348I) has recently been reported to reduce susceptibility to etravirine in subtypes A, B and C [32]. One clinical trial in South Africa found the N348I mutation present in 24% of patients failing first-line NNRTI regimens with subtype C virus, most commonly with nevirapine [33*]. This mutation is not included in standard mutation lists or algorithms but more data are urgently required to determine clinical relevance.

Rilpivirine (RPV) is another second-generation NNRTI with equal efficacy and similar patterns of resistance across subtypes [34,35]. However, regardless of subtype, RPV has suboptimal efficacy compared to efavirenz in ART-naïve individuals with HIV RNA >100 000 copies/ml [35].

Protease inhibitors

Non-polymorphic mutations in the protease gene have a greater impact on baseline susceptibility to protease inhibitors than polymorphic mutations [15]. However, recent evidence has suggested that the polymorphism at codon 36 in the protease gene (M36 in subtype B and I36 in most other subtypes) affects both the patterns of resistance that emerge under drug pressure and viral replication capacity [36]. Similarly, the M89 polymorphism in subtypes A, C, and CRF01_AE (L89 in subtype B) preferentially leads to the emergence under drug pressure of the M89T mutation, which confers high-level resistance to nelfinavir, atazanavir and lopinavir [37]. There is also *in vitro* evidence that CRF2_AG viruses with the 17E/64M polymorphisms demonstrate hypersusceptibility to certain protease inhibitors (nelfinavir, atazanavir and indinavir) [38].

Mutations in the *gag* cleavage sites and *gag* matrix protein are known to contribute to protease inhibitor resistance. Polymorphisms in this region may be more common in non-B subtypes [39,40] and have been shown to affect virological outcomes with lopinavir/ritonavir

monotherapy [41]. The importance of the subtype differences in *gag* are not well defined but may be more important in boosted PI monotherapy, which is under investigation for second-line therapy in resource-limited settings [42].

Integrase inhibitors

Raltegravir, a first-generation integrase inhibitor (INI), has demonstrated good efficacy in ART-naïve and ART-experienced individuals infected with different HIV-1 subtypes [43]. However, both raltegravir and the other first-generation INI elvitegravir have a relatively low genetic barrier to resistance. The primary mutations in the integrase gene associated with INI resistance are E92Q, Y143R/C, Q148K/R/H and N155H. The residues associated with primary resistance seem to be highly conserved across subtypes, but polymorphisms at the sites of secondary mutations are more common in non-B subtypes [8,44–46]. There is some evidence that the

effect of certain integrase mutations might differ according to subtype. Subtype B integrase enzyme with the N155H mutation (\pm E92Q) exhibited increased resistance to raltegravir compared to the subtype C enzyme [47].

Dolutegravir and MK-2048 are second-generation integrase inhibitors in development that have higher genetic barriers to resistance and retain activity against viruses with resistance to raltegravir or elvitegravir. There is some early evidence to suggest subtype differences could modulate the emergence of resistance to these drugs. In both INI naïve and raltegravir-experienced individuals, polymorphisms at codons 101 and 124 were more frequent in non-B subtypes than subtype B; these mutations were particularly prevalent in subtypes C and CRF02_AG [48]. Whilst the R263K mutation seems to be the most common mutation selected during dolutegravir therapy in subtype B, the G118R mutation previously associated

Table 2

Virological outcomes with specific antiretroviral drugs across different HIV-1 subtypes. Virological outcomes reported only for regimen including investigational drug, except for ARTEMIS where lopinavir and darunavir both reported

Clinical trials					
Drug name	Clinical trial(s)	Patient population	Participants (by subtype)	Virological outcomes ^a	Reference
Lopinavir	ARTEMIS	ART naïve	208 subtype B 50 subtype C 87 other subtypes	78% subtype B 82% subtype C 78% other subtypes	Dierynck 2010 [53]
Darunavir	ARTEMIS	ART naïve	210 subtype B 39 subtype C 93 other subtypes	81% subtype B 87% subtype C 88% other subtypes	Dierynck 2010 [53]
Etravirine	DUET-1, DUET-2	ART experienced	561 subtype B 33 non-B subtypes (no subtype C)	60% subtype B 73% non-B subtypes	Vingerhouts 2010 [28]
Rilpivirine	ECHO, THRIVE	ART naïve	485 subtype B 76 subtype C 125 other subtypes	84% subtype B 86% subtype C 86% other subtypes	Cohen 2012 [34]
Maraviroc	MERIT	ART naïve	Not reported	70% subtype B 61% subtype C 57% other subtypes	Cooper 2010 [54]
Raltegravir	STARTMRK	ART naïve	219 subtype B 19 subtype C 32 other subtypes	89% subtype B ^b 95% non-B subtypes ^b	Rockstroh 2011 [43]
Raltegravir	BENCHMRK-1, BENCHMRK-2	ART experienced	416 subtype B 3 subtype C 33 other subtypes	61% subtype B ^b 67% non-B subtypes ^b	Rockstroh 2011 [43]
Cohort studies					
Cohort	Drug regimens	Patient population	Participants (by subtype)	Virological outcomes	Reference
UK CHIC	2 NRTI + NNRTI 2 NRTI + boosted PI	ART naïve	1550 subtype B 272 subtype C 66 subtype A	89% subtype B ^c 94% subtype C ^c 97% subtype A ^c	Geretti 2009 [51]
Swiss HIV cohort	Various	ART naïve and mono/dual-NRTI experienced	2166 subtype B 383 non-B subtypes	89% subtype B ^d 90% non-B subtypes ^d	Scherrer 2011 [52*]

^a Outcome HIV RNA <50 copies/ml at week 48 unless otherwise stated.

^b HIV RNA <50 copies/ml at week 96; data not reported separately for subtype C.

^c HIV RNA <50 copies/ml at 12 months.

^d HIV RNA <50 copies/ml 90–365 days after ART initiation; results only shown for those who started ART 1999–2009.

with MK-2048 resistance might be a more common pathway in subtype C [49,50].

Clinical efficacy of new antiretroviral agents by HIV-1 subtype

Clinical evidence of different subtype responses to antiretroviral therapy (ART) might be the first indicator of subtype differences in the development of drug resistance. Inclusion of individuals infected with different subtypes is increasingly the norm in clinical trials of new antiretroviral agents, although the numbers infected with some non-B subtypes are quite low. Table 2 shows the virological responses by HIV-1 subtype in recently published clinical trials of new antiretroviral agents.

Cohort studies can also provide evidence of subtype differences in ART responses, although this is complicated by the use of different ART regimens. A collaborative group in the UK found that virological outcomes were broadly similar between subtypes A, B and C [51], while a more recent study from the Swiss HIV Cohort Study, which restricted analysis to white Caucasians, reported that those infected with non-B subtypes, had a lower risk of virological failure which was particularly apparent for subtypes A and CRF02_AG [52].

Conclusions

There is no compelling evidence that HIV-1 subtype needs be considered in the choice of ART regimens for first-line or second-line therapy, and other considerations of cost, effectiveness, toxicities and tolerability are more important in low-income and middle-income countries. However, recent evidence of subtype differences in drug resistance could potentially impact on antiretroviral strategies. The large amount of resistance data produced as part of surveillance studies and clinical care can be used to explore the differences in drug resistance between HIV-1 subtypes. National HIV drug resistance databases such as the ones in the UK, Switzerland and, recently, the Southern African Treatment and Resistance Network (SATuRN) Stanford and Rega public drug resistance databases are very useful national strategic resources to tackle the spread of drug resistance. The appreciation of subtype differences is also important to the development of new drugs, treatment strategies, drug sequencing, assessing response to treatment and surveillance for the transmission of resistance. In each of these areas, and in tracking the evolution of the HIV-1 pandemic, differences among subtypes continue to play an important role.

Acknowledgements

RJL is supported by the Wellcome Trust [grant number 090999]. TdO is supported by the Wellcome Trust [grant number 082384] and the grant entitled "Swiss-Prot/South Africa: Protein Bioinformatics Resource Development for Important Health-related Pathogens" under the Switzerland-South Africa Collaborative Research Program. European Commission (SANTE 2007 147-790).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Joint United Nations Programme on HIV/AIDS (UNAIDS): *World AIDS Day Report 2011*. Geneva, Switzerland: UNAIDS; 2011.
 2. Hamers RL, Kityo C, Lange JM, Wit TF, Mugenyi P: **Global threat from drug resistant HIV in sub-Saharan Africa**. *BMJ* 2012, **344**:e4159.
 3. Hemelaar J, Gouws E, Ghys PD, Osmanov S: **Global trends in molecular epidemiology of HIV-1 during 2000-2007**. *AIDS* 2011, **25**:679-689.
 4. Sharp PM, Hahn BH: **The evolution of HIV-1 and the origin of AIDS**. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences* 2010, **365**:2487-2494.
 5. Tebit DM, Arts EJ: **Tracking a century of global expansion and evolution of HIV to drive understanding and to combat disease**. *Lancet Infectious Diseases* 2011, **11**:45-56.
- Excellent overview of HIV-1 evolution, with particular focus on recombination.
6. Peeters M, Aghokeng AF, Delaporte E: **Genetic diversity among human immunodeficiency virus-1 non-B subtypes in viral load and drug resistance assays**. *Clinical Microbiology and Infection: The Official Publication of the European Society of Clinical Microbiology and Infectious Diseases* 2010, **16**:1525-1531.
 7. Hemelaar J: **The origin and diversity of the HIV-1 pandemic**. *Trends in Molecular Medicine* 2012, **18**:182-192.
 8. Brenner BG, Lowe M, Moisi D, Hardy I, Gagnon S, Charest H, Baril JG, Wainberg MA, Roger M: **Subtype diversity associated with the development of HIV-1 resistance to integrase inhibitors**. *Journal of Medical Virology* 2011, **83**:751-759.
 9. Chilton DN, Castro H, Lattimore S, Harrison LJ, Fearnhill E, Delpech V, Rice B, Pillay D, Dunn DT: **HIV type-1 drug resistance in antiretroviral treatment-naive adults infected with non-B subtype virus in the United Kingdom**. *Antiviral Therapy* 2010, **15**:985-991.
 10. Fox J, Castro H, Kaye S, McClure M, Weber JN, Fidler S: **Epidemiology of non-B clade forms of HIV-1 in men who have sex with men in the UK**. *AIDS* 2010, **24**:2397-2401.
 11. Wainberg MA, Zaharatos GJ, Brenner BG: **Development of antiretroviral drug resistance**. *New England Journal of Medicine* 2011, **365**:637-646.
 12. Kosakovsky Pond SL, Posada D, Stawiski E, Chappey C, Poon AF, Hughes G, Fearnhill E, Gravenor MB, Leigh Brown AJ, Frost SD: **An evolutionary model-based algorithm for accurate phylogenetic breakpoint mapping and subtype prediction in HIV-1**. *PLoS Computational Biology* 2009, **5**:e1000581.
 13. de Oliveira T, Shafer RW, Seebregts C: **Public database for HIV drug resistance in southern Africa**. *Nature* 2010, **464**:673.
 14. Melikian GL, Rhee SY, Taylor J, Fessel WJ, Kaufman D, Towner W, Troia-Cancio PV, Zolopa A, Robbins GK, Kagan R *et al.*: **Standardized comparison of the relative impacts of HIV-1 reverse transcriptase (RT) mutations on nucleoside RT inhibitor susceptibility**. *Antimicrobial Agents and Chemotherapy* 2012, **56**:2305-2313.
 15. Rhee SY, Taylor J, Fessel WJ, Kaufman D, Towner W, Troia P, Ruane P, Hellinger J, Shirvani V, Zolopa A *et al.*: **HIV-1 protease mutations and protease inhibitor cross-resistance**. *Antimicrobial Agents and Chemotherapy* 2010, **54**:4253-4261.
 16. Coutsinos D, Invernizzi CF, Xu H, Brenner BG, Wainberg MA: **Factors affecting template usage in the development of K65R resistance in subtype C variants of HIV type-1**. *Antiviral Chemistry and Chemotherapy* 2010, **20**:117-131.
 17. Coutsinos D, Invernizzi CF, Moisi D, Oliveira M, Martinez-Cajas JL, Brenner BG, Wainberg MA: **A template-dependent dislocation mechanism potentiates K65R reverse transcriptase mutation**

- development in subtype C variants of HIV-1.** *PLoS ONE* 2011, **6**:e20208.
Provides clear and detailed explanation for emergence of K65R in subtype C.
18. Kozal MJ, Chiarella J, St John EP, Moreno EA, Simen BB, Arnold TE, Lataillade M: **Prevalence of low-level HIV-1 variants with reverse transcriptase mutation K65R and the effect of antiretroviral drug exposure on variant levels.** *Antiviral Therapy* 2011, **16**:925-929.
 19. Li JF, Lipscomb JT, Wei X, Martinson NA, Morris L, Heneine W, Johnson JA: **Detection of low-level K65R variants in nucleoside reverse transcriptase inhibitor-naive chronic and acute HIV-1 subtype C infections.** *Journal of Infectious Diseases* 2011, **203**:798-802.
 20. Toni TA, Brenner BG, Asahchop EL, Ntemgwa M, Moisi D, Wainberg MA: **Development of an allele-specific PCR for detection of the K65R resistance mutation in patients infected with subtype C human immunodeficiency virus type 1.** *Antimicrobial Agents and Chemotherapy* 2010, **54**:907-911.
 21. Recordon-Pinson P, Papuchon J, Reigadas S, Deshpande A, Fleury H: **K65R in subtype C HIV-1 isolates from patients failing on a first-line regimen including d4T or AZT: comparison of Sanger and UDP sequencing data.** *PLoS ONE* 2012, **7**:e36549.
 22. Varghese V, Wang E, Babrzadeh F, Bachmann MH, Shahriar R, Liu T, Mappala SJ, Gharizadeh B, Fessel WJ, Katzenstein D *et al.*: **Nucleic acid template and the risk of a PCR-induced HIV-1 drug resistance mutation.** *PLoS ONE* 2010, **5**:e10992.
 23. Sunpath H, Wu B, Gordon M, Hampton J, Johnson B, Moosa MY, Ordonez C, Kuritzkes DR, Marconi VC: **High rate of K65R for ART naive patients with subtype C HIV infection failing a TDF-containing first-line regimen in South Africa.** *AIDS* 2012, **26**:1679-1684.
Concerning report from South Africa of very early development of K65R associated with virological failure of first-line ART regimens.
 24. Tang MW, Kanki PJ, Shafer RW: **A review of the virological efficacy of the 4 world health organization-recommended tenofovir-containing regimens for initial HIV therapy.** *Clinical Infectious Diseases* 2012, **54**:862-875.
A systematic review which finds evidence that regimens containing tenofovir and nevirapine might have suboptimal potency.
 25. Bansal V, Metzner KJ, Niederost B, Leemann C, Boni J, Gunthard HF, Fehr JS: **Minority K65R variants and early failure of antiretroviral therapy in HIV-1-infected Eritrean immigrant.** *Emerging Infectious Diseases* 2011, **17**:1966-1968.
 26. Eshleman SH, Hoover DR, Chen S, Hudelson SE, Guay LA, Mwatha A, Fiscus SA, Mmiro F, Musoke P, Jackson JB *et al.*: **Nevirapine (NVP) resistance in women with HIV-1 subtype C, compared with subtypes A and D, after the administration of single-dose NVP.** *Journal of Infectious Diseases* 2005, **192**:30-36.
 27. Kantor R: **Impact of HIV-1 pol diversity on drug resistance and its clinical implications.** *Current Opinion in Infectious Diseases* 2006, **19**:594-606.
 28. Vingerhoets J, Azijn H, Tambuyzer L, Dierynck I, De Meyer S, Rimsky L, Nijs S, De Smedt G, de Bethune MP, Picchio G: **Short communication: activity of etravirine on different HIV type 1 subtypes: in vitro susceptibility in treatment-naive patients and week 48 pooled DUET study data.** *AIDS Research and Human Retroviruses* 2010, **26**:621-624.
 29. Maiga AI, Descamps D, Morand-Joubert L, Malet I, Derache A, Cisse M, Koita V, Akonde A, Diarra B, Wiriden M *et al.*: **Resistance-associated mutations to etravirine (TMC-125) in antiretroviral-naive patients infected with non-B HIV-1 subtypes.** *Antimicrobial Agents and Chemotherapy* 2010, **54**:728-733.
 30. Asahchop EL, Oliveira M, Wainberg MA, Brenner BG, Moisi D, Toni T, Tremblay CL: **Characterization of the E138K resistance mutation in HIV-1 reverse transcriptase conferring susceptibility to etravirine in B and non-B HIV-1 subtypes.** *Antimicrobial Agents and Chemotherapy* 2011, **55**:600-607.
 31. Lai MT, Lu M, Felock PJ, Hrin RC, Wang YJ, Yan Y, Munshi S, McGaughey GB, Tynebor RM, Tucker TJ *et al.*: **Distinct mutation pathways of non-subtype B HIV-1 during in vitro resistance selection with nonnucleoside reverse transcriptase inhibitors.** *Antimicrobial Agents and Chemotherapy* 2010, **54**:4812-4824.
 32. McCormick AL, Parry CM, Crombe A, Goodall RL, Gupta RK, Kaleebu P, Kityo C, Chirara M, Towers GJ, Pillay D: **Impact of the N348I mutation in HIV-1 reverse transcriptase on nonnucleoside reverse transcriptase inhibitor resistance in non-subtype B HIV-1.** *Antimicrobial Agents and Chemotherapy* 2011, **55**:1806-1809.
 33. Brehm JH, Koontz DL, Wallis CL, Shutt KA, Sanne I, Wood R, McIntyre JA, Stevens WS, Sluis-Cremer N, Mellors JW: **Frequent emergence of N348I in HIV-1 subtype C reverse transcriptase with failure of initial therapy reduces susceptibility to reverse-transcriptase inhibitors.** *Clinical Infectious Diseases* 2012, **55**:737-745.
N348I is a mutation in the connection domain of reverse transcriptase not normally covered by standard genotyping methods. Here it was found frequently in patients failing first-line NNRTI-based therapy and was associated with reduced susceptibility to etravirine.
 34. Cohen CJ, Molina JM, Cahn P, Clotet B, Fourie J, Grinsztejn B, Wu H, Johnson MA, Saag M, Supparatpinyo K *et al.*: **Efficacy and safety of rilpivirine (TMC278) versus efavirenz at 48 weeks in treatment-naive HIV-1-infected patients: pooled results from the phase 3 double-blind randomized ECHO and THRIVE Trials.** *Journal of Acquired Immune Deficiency Syndromes* 2012, **60**:33-42.
 35. Rimsky L, Vingerhoets J, Van Eygen V, Eron J, Clotet B, Hoogstoel A, Boven K, Picchio G: **Genotypic and phenotypic characterization of HIV-1 isolates obtained from patients on rilpivirine therapy experiencing virologic failure in the phase 3 ECHO and THRIVE studies: 48-week analysis.** *Journal of Acquired Immune Deficiency Syndromes* 2012, **59**:39-46.
 36. Lisovsky I, Schader SM, Martinez-Cajas JL, Oliveira M, Moisi D, Wainberg MA: **HIV-1 protease codon 36 polymorphisms and differential development of resistance to nelfinavir, lopinavir, and atazanavir in different HIV-1 subtypes.** *Antimicrobial Agents and Chemotherapy* 2010, **54**:2878-2885.
 37. Martinez-Cajas JL, Wainberg MA, Oliveira M, Asahchop EL, Doualla-Bell F, Lisovsky I, Moisi D, Mendelson E, Grossman Z, Brenner BG: **The role of polymorphisms at position 89 in the HIV-1 protease gene in the development of drug resistance to HIV-1 protease inhibitors.** *Journal of Antimicrobial Chemotherapy* 2012, **67**:988-994.
 38. Santos AF, Tebit DM, Lalonde MS, Abecasis AB, Ratcliff A, Camacho RJ, Diaz RS, Herchenroder O, Soares MA, Arts EJ: **Effect of natural polymorphisms in the HIV-1 CRF02_AG protease on protease inhibitor hypersusceptibility.** *Antimicrobial Agents and Chemotherapy* 2012, **56**:2719-2725.
 39. Doyon L, Croteau G, Thibeault D, Poulin F, Pilote L, Lamarre D: **Second locus involved in human immunodeficiency virus type 1 resistance to protease inhibitors.** *Journal of Virology* 1996, **70**:3763-3769.
 40. Parry CM, Kolli M, Myers RE, Cane PA, Schiffer C, Pillay D: **Three residues in HIV-1 matrix contribute to protease inhibitor susceptibility and replication capacity.** *Antimicrobial Agents and Chemotherapy* 2011, **55**:1106-1113.
 41. Ghosn J, Delaugerre C, Flandre P, Galimand J, Cohen-Codar I, Raffi F, Delfraissy JF, Rouzioux C, Chaix ML: **Polymorphism in Gag gene cleavage sites of HIV-1 non-B subtype and virological outcome of a first-line lopinavir/ritonavir single drug regimen.** *PLoS ONE* 2011, **6**:e24798.
 42. Bartlett JA, Ribaldo HJ, Wallis CL, Aga E, Katzenstein DA, Stevens WS, Norton MR, Klingman KL, Hosseinipour MC, Crump JA *et al.*: **Lopinavir/ritonavir monotherapy after virologic failure of first-line antiretroviral therapy in resource-limited settings.** *AIDS* 2012, **26**:1345-1354.
 43. Rockstroh JK, Teppler H, Zhao J, Sklar P, Miller MD, Harvey CM, Strohmaier KM, Leavitt RY, Nguyen BY: **Clinical efficacy of raltegravir against B and non-B subtype HIV-1 in phase III clinical studies.** *AIDS* 2011, **25**:1365-1369.
 44. Fish MQ, Hewer R, Wallis CL, Venter WD, Stevens WS, Papathanasopoulos MA: **Natural polymorphisms of integrase**

- among HIV type 1-infected South African patients. *AIDS Research and Human Retroviruses* 2010, **26**:489-493.
45. Garrido C, Geretti AM, Zahonero N, Booth C, Strang A, Soriano V, De Mendoza C: **Integrase variability and susceptibility to HIV integrase inhibitors: impact of subtypes, antiretroviral experience and duration of HIV infection.** *Journal of Antimicrobial Chemotherapy* 2010, **65**:320-326.
 46. Sierra S, Lubke N, Walter H, Schuler E, Reuter S, Fatkenheuer G, Bickel M, da Silva H, Kaiser R, Esser S: **The SnoB study: frequency of baseline raltegravir resistance mutations prevalence in different non-B subtypes.** *Medical Microbiology and Immunology* 2011, **200**:225-232.
 47. Bar-Magen T, Donahue DA, McDonough EI, Kuhl BD, Faltenbacher VH, Xu H, Michaud V, Sloan RD, Wainberg MA: **HIV-1 subtype B and C integrase enzymes exhibit differential patterns of resistance to integrase inhibitors in biochemical assays.** *AIDS* 2010, **24**:2171-2179.
 48. Garrido C, Soriano V, Geretti AM, Zahonero N, Garcia S, Booth C, Gutierrez F, Viciano I, de Mendoza C: **Resistance associated mutations to dolutegravir (S/GSK1349572) in HIV-infected patients – impact of HIV subtypes and prior raltegravir experience.** *Antiviral Research* 2011, **90**:164-167.
 49. Quashie PK, Mesplede T, Han YS, Oliveira M, Singhroy DN,
 - Fujiwara T, Underwood MR, Wainberg MA: **Characterization of the R263K mutation in HIV-1 integrase that confers low-level resistance to the second-generation integrase strand transfer inhibitor dolutegravir.** *Journal of Virology* 2012, **86**:2696-2705.
 This study suggests that the next-generation integrase inhibitor dolutegravir might have different resistance pathways in subtype B and C.
 50. Bar-Magen T, Sloan RD, Donahue DA, Kuhl BD, Zabeida A, Xu H, Oliveira M, Hazuda DJ, Wainberg MA: **Identification of novel mutations responsible for resistance to MK-2048, a second-generation HIV-1 integrase inhibitor.** *Journal of Virology* 2010, **84**:9210-9216.
 51. Geretti AM, Harrison L, Green H, Sabin C, Hill T, Fearnhill E, Pillay D, Dunn D: **Effect of HIV-1 subtype on virologic and immunologic response to starting highly active antiretroviral therapy.** *Clinical Infectious Diseases* 2009, **48**:1296-1305.
 52. Scherrer AU, Ledergerber B, von Wyl V, Boni J, Yerly S, Klimkait T,
 - Burgisser P, Rauch A, Hirschel B, Cavassini M *et al.*: **Improved virological outcome in White patients infected with HIV-1 non-B subtypes compared to subtype B.** *Clinical Infectious Diseases* 2011, **53**:1143-1152.
 This large cohort study was able to demonstrate that virological outcomes in non-B subtypes are at least as good as, if not better than, subtype B.
 53. Dierynck I, De Meyer S, Lathouwers E, Vanden Abeele C, Van De Castele T, Spinosa-Guzman S, de Bethune MP, Picchio G: **In vitro susceptibility and virological outcome to darunavir and lopinavir are independent of HIV type-1 subtype in treatment-naive patients.** *Antiviral Therapy* 2010, **15**:1161-1169.
 54. Cooper DA, Heera J, Goodrich J, Tawadrous M, Saag M, Dejesus E, Clumeck N, Walmsley S, Ting N, Coakley E *et al.*: **Maraviroc versus efavirenz, both in combination with zidovudine-lamivudine, for the treatment of antiretroviral-naive subjects with CCR5-tropic HIV-1 infection.** *Journal of Infectious Diseases* 2010, **201**:803-813.